

# A dynamical system related to GIT

Nolan R. Wallach

June 4, 2015

## A gradient system

- Let  $\phi \in \mathbb{R}[x_1, \dots, x_n]$  be a polynomial that is homogeneous of degree  $m$  such that  $\phi(x) \geq 0$  for all  $x \in \mathbb{R}^n$ . We consider the gradient system

$$\frac{dx}{dt} = -\nabla\phi(x)$$

## A gradient system

- Let  $\phi \in \mathbb{R}[x_1, \dots, x_n]$  be a polynomial that is homogeneous of degree  $m$  such that  $\phi(x) \geq 0$  for all  $x \in \mathbb{R}^n$ . We consider the gradient system

$$\frac{dx}{dt} = -\nabla\phi(x)$$

- Note that

$$\langle \nabla\phi(x), x \rangle = m\phi(x)$$

Denoting by  $F(t, x)$  the solution to the system near  $t = 0$  with  $F(0, x) = x$ . Then

$$\begin{aligned} \frac{d}{dt} \langle F(t, x), F(t, x) \rangle &= -2 \langle \nabla\phi(F(t, x)), F(t, x) \rangle \\ &= -2m\phi(F(t, x)) \leq 0. \end{aligned}$$

- This implies  $\|F(t, x)\| \leq \|x\|$  where defined for  $t \geq 0$  and hence  $F(t, x)$  is defined for all  $t \geq 0$ .

- This implies  $\|F(t, x)\| \leq \|x\|$  where defined for  $t \geq 0$  and hence  $F(t, x)$  is defined for all  $t \geq 0$ .
- The formula

$$\langle \nabla \phi(x), x \rangle = m\phi(x)$$

combined with the Schwarz inequality implies that

$$\|\nabla \phi(x)\| \|x\| \geq m\phi(x).$$

- This implies  $\|F(t, x)\| \leq \|x\|$  where defined for  $t \geq 0$  and hence  $F(t, x)$  is defined for all  $t \geq 0$ .
- The formula

$$\langle \nabla \phi(x), x \rangle = m\phi(x)$$

combined with the Schwarz inequality implies that

$$\|\nabla \phi(x)\| \|x\| \geq m\phi(x).$$

- The Lojasiewicz gradient inequality implies the following improvement. There exists  $0 < \varepsilon \leq \frac{1}{m-1}$  and  $C > 0$  both depending only on  $\phi$  such that

$$\|\nabla \phi(x)\|^{1+\varepsilon} \|x\|^{1-(m-1)\varepsilon} \geq C\phi(x).$$

- We take  $\varepsilon$  and  $C$  as above (but allow  $\varepsilon = 0$  which is easy). If we write  $F$  for  $F(t, X)$  and  $H(t) = \phi(F(t, x))$  then we have

$$H'(t) = -d\phi(F)\nabla\phi(F) = -\|\nabla\phi(F)\|^2.$$

- We take  $\varepsilon$  and  $C$  as above (but allow  $\varepsilon = 0$  which is easy). If we write  $F$  for  $F(t, X)$  and  $H(t) = \phi(F(t, x))$  then we have

$$H'(t) = -d\phi(F)\nabla\phi(F) = -\|\nabla\phi(F)\|^2.$$

- If  $t \geq 0$  and  $\|x\| \leq r$

$$\|\nabla\phi(F)\|^{1+\varepsilon} r^{1-(m-1)\varepsilon} \geq \|\nabla\phi(F)\|^{1+\varepsilon} \|F\|^{1-(m-1)\varepsilon} \geq C\phi(x).$$



- We take  $\varepsilon$  and  $C$  as above (but allow  $\varepsilon = 0$  which is easy). If we write  $F$  for  $F(t, X)$  and  $H(t) = \phi(F(t, x))$  then we have

$$H'(t) = -d\phi(F)\nabla\phi(F) = -\|\nabla\phi(F)\|^2.$$

- If  $t \geq 0$  and  $\|x\| \leq r$

$$\|\nabla\phi(F)\|^{1+\varepsilon} r^{1-(m-1)\varepsilon} \geq \|\nabla\phi(F)\|^{1+\varepsilon} \|F\|^{1-(m-1)\varepsilon} \geq C\phi(x).$$

- We will now run through what has come to be called “the Lojasiewicz argument” which I learned from a beautiful exposition of Neeman’s theorem by Gerry Schwarz.

$$\|\nabla\phi(F)\|^{1+\varepsilon} \geq \frac{C}{r^{1-3\varepsilon}}\phi(F).$$

- 

$$\|\nabla\phi(F)\|^{1+\varepsilon} \geq \frac{C}{r^{1-3\varepsilon}}\phi(F).$$

- 

$$\|\nabla\phi(F)\|^2 \geq \left(\frac{C}{r^{1-3\varepsilon}}\right)^{\frac{2}{1+\varepsilon}} \phi(F)^{\frac{2}{1+\varepsilon}}.$$

- $$\|\nabla\phi(F)\|^{1+\varepsilon} \geq \frac{C}{r^{1-3\varepsilon}}\phi(F).$$

- $$\|\nabla\phi(F)\|^2 \geq \left(\frac{C}{r^{1-3\varepsilon}}\right)^{\frac{2}{1+\varepsilon}} \phi(F)^{\frac{2}{1+\varepsilon}}.$$

- $$|H'(t)| \geq \frac{1}{2} \left(\frac{C}{r^{1-3\varepsilon}}\right)^{\frac{2}{1+\varepsilon}} \phi(F)^{\frac{2}{1+\varepsilon}} = C_1(r)H(t)^{\frac{2}{1+\varepsilon}}.$$

- $$\|\nabla\phi(F)\|^{1+\varepsilon} \geq \frac{C}{r^{1-3\varepsilon}}\phi(F).$$

- $$\|\nabla\phi(F)\|^2 \geq \left(\frac{C}{r^{1-3\varepsilon}}\right)^{\frac{2}{1+\varepsilon}} \phi(F)^{\frac{2}{1+\varepsilon}}.$$

- $$|H'(t)| \geq \frac{1}{2} \left(\frac{C}{r^{1-3\varepsilon}}\right)^{\frac{2}{1+\varepsilon}} \phi(F)^{\frac{2}{1+\varepsilon}} = C_1(r)H(t)^{\frac{2}{1+\varepsilon}}.$$

- Since  $H'(t) \leq 0$  for  $t \geq 0$  we have  $-H'(t) \geq C_1(r)H(t)^{\frac{2}{1+\varepsilon}}$ .  
 Assuming  $H(t) > 0$  we have

- $$\|\nabla\phi(F)\|^{1+\varepsilon} \geq \frac{C}{r^{1-3\varepsilon}}\phi(F).$$

- $$\|\nabla\phi(F)\|^2 \geq \left(\frac{C}{r^{1-3\varepsilon}}\right)^{\frac{2}{1+\varepsilon}} \phi(F)^{\frac{2}{1+\varepsilon}}.$$

- $$|H'(t)| \geq \frac{1}{2} \left(\frac{C}{r^{1-3\varepsilon}}\right)^{\frac{2}{1+\varepsilon}} \phi(F)^{\frac{2}{1+\varepsilon}} = C_1(r)H(t)^{\frac{2}{1+\varepsilon}}.$$

- Since  $H'(t) \leq 0$  for  $t \geq 0$  we have  $-H'(t) \geq C_1(r)H(t)^{\frac{2}{1+\varepsilon}}$ .  
Assuming  $H(t) > 0$  we have

- $$\frac{d}{dt}H(t)^{-\frac{1-\varepsilon}{1+\varepsilon}} = -\frac{1-\varepsilon}{1+\varepsilon} \frac{H'(t)}{H(t)^{\frac{2}{1+\varepsilon}}} \geq C_1(r)$$

$$H(t)^{-\frac{1-\epsilon}{1+\epsilon}} \geq C_1(r)t.$$

- 

$$H(t)^{-\frac{1-\varepsilon}{1+\varepsilon}} \geq C_1(r)t.$$

- 

$$H(t) \leq C_2(r)t^{-\frac{(1+\varepsilon)}{1-\varepsilon}} \leq C_2(r)t^{-(1+\varepsilon)},$$





$$H(t)^{-\frac{1-\varepsilon}{1+\varepsilon}} \geq C_1(r)t.$$



$$H(t) \leq C_2(r)t^{-\frac{(1+\varepsilon)}{1-\varepsilon}} \leq C_2(r)t^{-(1+\varepsilon)},$$

- This is true if  $H(t) = 0$  so the formula is valid for all  $t > 0$ .



$$H(t)^{-\frac{1-\varepsilon}{1+\varepsilon}} \geq C_1(r)t.$$



$$H(t) \leq C_2(r)t^{-\frac{(1+\varepsilon)}{1-\varepsilon}} \leq C_2(r)t^{-(1+\varepsilon)},$$

- This is true if  $H(t) = 0$  so the formula is valid for all  $t > 0$ .
- This is the first half of the calculus part of the Lojasiewicz argument. The first implication needs only the easy case  $\varepsilon = 0$ . If  $\|x\| \leq r$  then

$$\phi(F(t, x)) \leq \frac{C(r)}{t}$$

so  $\lim_{t \rightarrow +\infty} \phi(F(t, x)) = 0$  uniformly for  $x$  in compacta. We now do the rest of the Lojasiewicz argument which uses the existence of  $\varepsilon > 0$ .

- Let  $f(t) = t^{1+\delta}$  with  $0 < \delta < \varepsilon$  then for  $t > 0$

$$0 < H(t)f'(t) \leq C_2(r)(1 + \delta)t^{-1-(\varepsilon-\delta)}.$$

- Let  $f(t) = t^{1+\delta}$  with  $0 < \delta < \varepsilon$  then for  $t > 0$

$$0 < H(t)f'(t) \leq C_2(r)(1 + \delta)t^{-1-(\varepsilon-\delta)}.$$



$$\begin{aligned} H(s)f(s) - H(t)f(t) &= \int_t^s \frac{d}{du}(H(u)f(u)) du = \\ &= \int_t^s H(u)f'(u) du + \int_t^s H'(u)f(u) du. \end{aligned}$$

- Let  $f(t) = t^{1+\delta}$  with  $0 < \delta < \varepsilon$  then for  $t > 0$

$$0 < H(t)f'(t) \leq C_2(r)(1+\delta)t^{-1-(\varepsilon-\delta)}.$$

- $$H(s)f(s) - H(t)f(t) = \int_t^s \frac{d}{du}(H(u)f(u)) du =$$
$$\int_t^s H(u)f'(u) du + \int_t^s H'(u)f(u) du.$$

- $$- \int_t^s H'(u)f(u) du = \int_t^s H(u)f'(u) du + H(t)f(t) - H(s)f(s).$$

$$0 \leq H(s)f(s) \leq C_2(r)s^{-(1+\varepsilon)}s^{1+\delta} = C_2(r)s^{-(\varepsilon-\delta)}.$$

- Let  $f(t) = t^{1+\delta}$  with  $0 < \delta < \varepsilon$  then for  $t > 0$

$$0 < H(t)f'(t) \leq C_2(r)(1+\delta)t^{-1-(\varepsilon-\delta)}.$$

- 

$$H(s)f(s) - H(t)f(t) = \int_t^s \frac{d}{du}(H(u)f(u)) du =$$

$$\int_t^s H(u)f'(u) du + \int_t^s H'(u)f(u) du.$$

- 

$$- \int_t^s H'(u)f(u) du = \int_t^s H(u)f'(u) du + H(t)f(t) - H(s)f(s).$$

$$0 \leq H(s)f(s) \leq C_2(r)s^{-(1+\varepsilon)}s^{1+\delta} = C_2(r)s^{-(\varepsilon-\delta)}.$$

- 

$$\lim_{s \rightarrow +\infty} \int_t^s |H'(u)| f(u) du = \int_t^\infty H(u)f'(u) du + H(t)f(t).$$

- Thus  $\sqrt{|H'(u)| f(u)}$  is in  $L^2([t, +\infty))$  for all  $t > 0$  and so

$$\sqrt{|H'(u)|} = \sqrt{|H'(u)| f(u) u^{-\frac{(1+\delta)}{2}}} \in L^1([t, +\infty)).$$

- Thus  $\sqrt{|H'(u)| f(u)}$  is in  $L^2([t, +\infty))$  for all  $t > 0$  and so

$$\sqrt{|H'(u)|} = \sqrt{|H'(u)| f(u) u^{-\frac{(1+\delta)}{2}}} \in L^1([t, +\infty)).$$

- **Theorem.** If  $t > 0$  then

$$\int_t^{+\infty} \left\| \frac{d}{du} F(u, x) \right\| du$$

converges uniformly for  $\|x\| \leq r$ .



- Thus  $\sqrt{|H'(u)| f(u)}$  is in  $L^2([t, +\infty))$  for all  $t > 0$  and so

$$\sqrt{|H'(u)|} = \sqrt{|H'(u)| f(u) u^{-\frac{(1+\delta)}{2}}} \in L^1([t, +\infty)).$$

- **Theorem.** If  $t > 0$  then

$$\int_t^{+\infty} \left\| \frac{d}{du} F(u, x) \right\| du$$

converges uniformly for  $\|x\| \leq r$ .

- 

$$\int_t^{\infty} \frac{d}{du} F(u, x) du$$

converges absolutely and uniformly for  $\|x\| \leq r$ .

- Thus  $\sqrt{|H'(u)| f(u)}$  is in  $L^2([t, +\infty))$  for all  $t > 0$  and so

$$\sqrt{|H'(u)|} = \sqrt{|H'(u)| f(u) u^{-\frac{(1+\delta)}{2}}} \in L^1([t, +\infty)).$$

- **Theorem.** If  $t > 0$  then

$$\int_t^{+\infty} \left\| \frac{d}{du} F(u, x) \right\| du$$

converges uniformly for  $\|x\| \leq r$ .

- 

$$\int_t^{\infty} \frac{d}{du} F(u, x) du$$

converges absolutely and uniformly for  $\|x\| \leq r$ .

- Noting that if  $s > t$  then

$$\int_t^s \frac{d}{du} F(u, x) du = F(s, x) - F(t, x)$$

we have for  $t > 0$

$$\lim_{s \rightarrow \infty} F(s, x) = \int_t^{\infty} \frac{d}{du} F(u, x) du + F(t, x).$$

- Finally, set  $L(t, x) = F(\frac{t}{1-t}, x)$  and define  $L(1, x)$  by the limit above then  $L : [0, 1] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is continuous and since

$$\nabla \phi(x) = 0 \iff \phi(x) = 0$$

we have

- Finally, set  $L(t, x) = F(\frac{t}{1-t}, x)$  and define  $L(1, x)$  by the limit above then  $L : [0, 1] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is continuous and since

$$\nabla\phi(x) = 0 \iff \phi(x) = 0$$

we have

- **Theorem.**  $L : [0, 1] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  defines a strong deformation retraction of  $\mathbb{R}^n$  onto  $Y = \{x \in \mathbb{R}^n | \phi(x) = 0\}$ .

- Finally, set  $L(t, x) = F(\frac{t}{1-t}, x)$  and define  $L(1, x)$  by the limit above then  $L : [0, 1] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is continuous and since

$$\nabla\phi(x) = 0 \iff \phi(x) = 0$$

we have

- **Theorem.**  $L : [0, 1] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  defines a strong deformation retraction of  $\mathbb{R}^n$  onto  $Y = \{x \in \mathbb{R}^n | \phi(x) = 0\}$ .
- **Corollary.** If  $Z \subset \mathbb{R}^n$  is closed and such that  $F(t, z) \in Z$  for  $t \geq 0$  and  $z \in Z$  then  $H : [0, 1] \times Z \rightarrow Z$  defines a strong deformation retraction of  $Z$  onto  $Z \cap Y$ .

## Kempf-Ness over the reals

- Let  $G$  be an open subgroup of a Zariski closed subgroup of  $GL(n, \mathbb{R})$  that is closed under real adjoint relative to the standard inner product,  $\langle \dots, \dots \rangle$ ,  $g \rightarrow g^*$ . Let  $K = G \cap O(n)$ . Then  $K$  is a maximal compact subgroup of  $G$ . On  $\mathfrak{g} = Lie(G)$  we put the inner product  $\langle X, Y \rangle = tr(XY^*)$ , Set  $\mathfrak{p} = Lie(K)^\perp$  relative to this inner product.

## Kempf-Ness over the reals

- Let  $G$  be an open subgroup of a Zariski closed subgroup of  $GL(n, \mathbb{R})$  that is closed under real adjoint relative to the standard inner product,  $\langle \dots, \dots \rangle$ ,  $g \rightarrow g^*$ . Let  $K = G \cap O(n)$ . Then  $K$  is a maximal compact subgroup of  $G$ . On  $\mathfrak{g} = Lie(G)$  we put the inner product  $\langle X, Y \rangle = tr(XY^*)$ , Set  $\mathfrak{p} = Lie(K)^\perp$  relative to this inner product.
- We say that an element  $v \in \mathbb{R}^n$  is  $G$ -critical if for any  $X \in Lie(G)$ ,  $\langle Xv, v \rangle = 0$ . The following is an extension of the Kempf-Ness Theorem first observed by Richardson and Slodoway.

## Kempf-Ness over the reals

- Let  $G$  be an open subgroup of a Zariski closed subgroup of  $GL(n, \mathbb{R})$  that is closed under real adjoint relative to the standard inner product,  $\langle \dots, \dots \rangle$ ,  $g \rightarrow g^*$ . Let  $K = G \cap O(n)$ . Then  $K$  is a maximal compact subgroup of  $G$ . On  $\mathfrak{g} = Lie(G)$  we put the inner product  $\langle X, Y \rangle = tr(XY^*)$ , Set  $\mathfrak{p} = Lie(K)^\perp$  relative to this inner product.
- We say that an element  $v \in \mathbb{R}^n$  is  $G$ -critical if for any  $X \in Lie(G)$ ,  $\langle Xv, v \rangle = 0$ . The following is an extension of the Kempf-Ness Theorem first observed by Richardson and Slodoway.
- **Theorem.** Let  $G, K$  be as above. Let  $v \in \mathbb{R}^n$ .



## Kempf-Ness over the reals

- Let  $G$  be an open subgroup of a Zariski closed subgroup of  $GL(n, \mathbb{R})$  that is closed under real adjoint relative to the standard inner product,  $\langle \dots, \dots \rangle$ ,  $g \rightarrow g^*$ . Let  $K = G \cap O(n)$ . Then  $K$  is a maximal compact subgroup of  $G$ . On  $\mathfrak{g} = Lie(G)$  we put the inner product  $\langle X, Y \rangle = tr(XY^*)$ , Set  $\mathfrak{p} = Lie(K)^\perp$  relative to this inner product.
- We say that an element  $v \in \mathbb{R}^n$  is  $G$ -critical if for any  $X \in Lie(G)$ ,  $\langle Xv, v \rangle = 0$ . The following is an extension of the Kempf-Ness Theorem first observed by Richardson and Slodoway.
- **Theorem.** Let  $G, K$  be as above. Let  $v \in \mathbb{R}^n$ .
- ① If  $v$  is critical if and only if  $\|gv\| \geq \|v\|$  for all  $g \in G$ .

## Kempf-Ness over the reals

- Let  $G$  be an open subgroup of a Zariski closed subgroup of  $GL(n, \mathbb{R})$  that is closed under real adjoint relative to the standard inner product,  $\langle \dots, \dots \rangle$ ,  $g \rightarrow g^*$ . Let  $K = G \cap O(n)$ . Then  $K$  is a maximal compact subgroup of  $G$ . On  $\mathfrak{g} = Lie(G)$  we put the inner product  $\langle X, Y \rangle = tr(XY^*)$ , Set  $\mathfrak{p} = Lie(K)^\perp$  relative to this inner product.
- We say that an element  $v \in \mathbb{R}^n$  is  $G$ -critical if for any  $X \in Lie(G)$ ,  $\langle Xv, v \rangle = 0$ . The following is an extension of the Kempf-Ness Theorem first observed by Richardson and Slodoway.
- **Theorem.** Let  $G, K$  be as above. Let  $v \in \mathbb{R}^n$ .
  - 1 If  $v$  is critical if and only if  $\|gv\| \geq \|v\|$  for all  $g \in G$ .
  - 2 If  $v$  is critical and if  $w \in Gv$  is such that  $\|v\| = \|w\|$  then  $w \in Kv$ .

## Kempf-Ness over the reals

- Let  $G$  be an open subgroup of a Zariski closed subgroup of  $GL(n, \mathbb{R})$  that is closed under real adjoint relative to the standard inner product,  $\langle \dots, \dots \rangle$ ,  $g \rightarrow g^*$ . Let  $K = G \cap O(n)$ . Then  $K$  is a maximal compact subgroup of  $G$ . On  $\mathfrak{g} = Lie(G)$  we put the inner product  $\langle X, Y \rangle = tr(XY^*)$ , Set  $\mathfrak{p} = Lie(K)^\perp$  relative to this inner product.
- We say that an element  $v \in \mathbb{R}^n$  is  $G$ -critical if for any  $X \in Lie(G)$ ,  $\langle Xv, v \rangle = 0$ . The following is an extension of the Kempf-Ness Theorem first observed by Richardson and Slodoway.
- **Theorem.** Let  $G, K$  be as above. Let  $v \in \mathbb{R}^n$ .
  - 1 If  $v$  is critical if and only if  $\|gv\| \geq \|v\|$  for all  $g \in G$ .
  - 2 If  $v$  is critical and if  $w \in Gv$  is such that  $\|v\| = \|w\|$  then  $w \in Kv$ .
  - 3 If  $Gv$  is closed then there exists a critical element in  $Gv$ .

## Kempf-Ness over the reals

- Let  $G$  be an open subgroup of a Zariski closed subgroup of  $GL(n, \mathbb{R})$  that is closed under real adjoint relative to the standard inner product,  $\langle \dots, \dots \rangle$ ,  $g \rightarrow g^*$ . Let  $K = G \cap O(n)$ . Then  $K$  is a maximal compact subgroup of  $G$ . On  $\mathfrak{g} = Lie(G)$  we put the inner product  $\langle X, Y \rangle = tr(XY^*)$ , Set  $\mathfrak{p} = Lie(K)^\perp$  relative to this inner product.
- We say that an element  $v \in \mathbb{R}^n$  is  $G$ -critical if for any  $X \in Lie(G)$ ,  $\langle Xv, v \rangle = 0$ . The following is an extension of the Kempf-Ness Theorem first observed by Richardson and Slodoway.
- **Theorem.** Let  $G, K$  be as above. Let  $v \in \mathbb{R}^n$ .
  - 1 If  $v$  is critical if and only if  $\|gv\| \geq \|v\|$  for all  $g \in G$ .
  - 2 If  $v$  is critical and if  $w \in Gv$  is such that  $\|v\| = \|w\|$  then  $w \in Kv$ .
  - 3 If  $Gv$  is closed then there exists a critical element in  $Gv$ .
  - 4 If  $v$  is critical then  $Gv$  is closed.

- We set  $V = \mathbb{R}^n$  as a  $G$ -module and  $Crit_G(V)$  equal to the set of all critical vectors. If  $X_1, \dots, X_r$  is an orthonormal basis of  $\mathfrak{p}$  then

$$\phi(v) = \sum \langle X_j v, v \rangle^2$$

is non-negative homogeneous polynomial of degree 4 defining  $Crit_G(V)$ .

- We set  $V = \mathbb{R}^n$  as a  $G$ -module and  $Crit_G(V)$  equal to the set of all critical vectors. If  $X_1, \dots, X_r$  is an orthonormal basis of  $\mathfrak{p}$  then

$$\phi(v) = \sum \langle X_j v, v \rangle^2$$

is non-negative homogeneous polynomial of degree 4 defining  $Crit_G(V)$ .

- We consider  $\mathbb{R}^n$  as  $n \times 1$  columns and thus if  $v \in V$  then  $v^*$  is  $v$  as a row vector. So for  $v, w \in V$ ,  $vw^*$  is an  $n \times n$  matrix and

$$\langle Xv, w \rangle = \text{tr}Xvw^*.$$

- We set  $V = \mathbb{R}^n$  as a  $G$ -module and  $Crit_G(V)$  equal to the set of all critical vectors. If  $X_1, \dots, X_r$  is an orthonormal basis of  $\mathfrak{p}$  then

$$\phi(v) = \sum \langle X_j v, v \rangle^2$$

is non-negative homogeneous polynomial of degree 4 defining  $Crit_G(V)$ .

- We consider  $\mathbb{R}^n$  as  $n \times 1$  columns and thus if  $v \in V$  then  $v^*$  is  $v$  as a row vector. So for  $v, w \in V$ ,  $vw^*$  is an  $n \times n$  matrix and

$$\langle Xv, w \rangle = \text{tr}Xvw^*.$$

- Let  $P_{\mathfrak{g}}$  be the orthogonal projection of  $M_n(\mathbb{R})$  onto  $\mathfrak{g}$  then

$$\nabla\phi(v) = 4P_{\mathfrak{g}}(vw^*)v \in T_v(Gv).$$

- We set  $V = \mathbb{R}^n$  as a  $G$ -module and  $Crit_G(V)$  equal to the set of all critical vectors. If  $X_1, \dots, X_r$  is an orthonormal basis of  $\mathfrak{p}$  then

$$\phi(v) = \sum \langle X_j v, v \rangle^2$$

is non-negative homogeneous polynomial of degree 4 defining  $Crit_G(V)$ .

- We consider  $\mathbb{R}^n$  as  $n \times 1$  columns and thus if  $v \in V$  then  $v^*$  is  $v$  as a row vector. So for  $v, w \in V$ ,  $vw^*$  is an  $n \times n$  matrix and

$$\langle Xv, w \rangle = \text{tr}Xvw^*.$$

- Let  $P_{\mathfrak{g}}$  be the orthogonal projection of  $M_n(\mathbb{R})$  onto  $\mathfrak{g}$  then

$$\nabla\phi(v) = 4P_{\mathfrak{g}}(vw^*)v \in T_v(Gv).$$

- Also note that  $\nabla\phi(kv) = k\nabla\phi(v)$  for  $k \in K$ .



- Let  $F(t, x)$  be the gradient flow corresponding to  $\phi$ . Then we have shown using freshman calculus that for  $t > 0$  and  $\|x\| \leq r$

$$\phi(F(t, x)) \leq \frac{C(r)}{t}.$$

- Let  $F(t, x)$  be the gradient flow corresponding to  $\phi$ . Then we have shown using freshman calculus that for  $t > 0$  and  $\|x\| \leq r$

$$\phi(F(t, x)) \leq \frac{C(r)}{t}.$$

- In addition if  $Z \subset V$  is closed and  $G$ -invariant then  $F(t, Z) \subset Z$  and 2 in the real Kempf-Ness theorem implies:

- Let  $F(t, x)$  be the gradient flow corresponding to  $\phi$ . Then we have shown using freshman calculus that for  $t > 0$  and  $\|x\| \leq r$

$$\phi(F(t, x)) \leq \frac{C(r)}{t}.$$

- In addition if  $Z \subset V$  is closed and  $G$ -invariant then  $F(t, Z) \subset Z$  and 2 in the real Kempf-Ness theorem implies:
- **Theorem.** Setting  $L(t, Kv) = KF(\frac{t}{1-t}, v)$   $0 \leq t < 1$  then  $\lim_{t \rightarrow 1} L(t, Kv)$  converges uniformly on compacta and this yields a strict deformation retraction of  $Z/K$  to  $(\text{Crit}_G(V) \cap Z) / K$  for any  $G$ -invariant closed subset of  $V$ .

- Let  $F(t, x)$  be the gradient flow corresponding to  $\phi$ . Then we have shown using freshman calculus that for  $t > 0$  and  $\|x\| \leq r$

$$\phi(F(t, x)) \leq \frac{C(r)}{t}.$$

- In addition if  $Z \subset V$  is closed and  $G$ -invariant then  $F(t, Z) \subset Z$  and 2 in the real Kempf-Ness theorem implies:
- **Theorem.** Setting  $L(t, Kv) = KF(\frac{t}{1-t}, v)$   $0 \leq t < 1$  then  $\lim_{t \rightarrow 1} L(t, Kv)$  converges uniformly on compacta and this yields a strict deformation retraction of  $Z/K$  to  $(\text{Crit}_G(V) \cap Z) / K$  for any  $G$ -invariant closed subset of  $V$ .
- The statement of the next result is simplified.

- Let  $F(t, x)$  be the gradient flow corresponding to  $\phi$ . Then we have shown using freshman calculus that for  $t > 0$  and  $\|x\| \leq r$

$$\phi(F(t, x)) \leq \frac{C(r)}{t}.$$

- In addition if  $Z \subset V$  is closed and  $G$ -invariant then  $F(t, Z) \subset Z$  and 2 in the real Kempf-Ness theorem implies:
- **Theorem.** Setting  $L(t, Kv) = KF(\frac{t}{1-t}, v)$   $0 \leq t < 1$  then  $\lim_{t \rightarrow 1} L(t, Kv)$  converges uniformly on compacta and this yields a strict deformation retraction of  $Z/K$  to  $(\text{Crit}_G(V) \cap Z) / K$  for any  $G$ -invariant closed subset of  $V$ .
- The statement of the next result is simplified.
- **Corollary.** If  $Z$  is Zariski closed in  $V$  then the GIT quotient,  $Z // G$ , of  $Z$  is a strict deformation retract of  $Z/K$ .

- Let  $F(t, x)$  be the gradient flow corresponding to  $\phi$ . Then we have shown using freshman calculus that for  $t > 0$  and  $\|x\| \leq r$

$$\phi(F(t, x)) \leq \frac{C(r)}{t}.$$

- In addition if  $Z \subset V$  is closed and  $G$ -invariant then  $F(t, Z) \subset Z$  and 2 in the real Kempf-Ness theorem implies:
- Theorem.** Setting  $L(t, Kv) = KF(\frac{t}{1-t}, v)$   $0 \leq t < 1$  then  $\lim_{t \rightarrow 1} L(t, Kv)$  converges uniformly on compacta and this yields a strict deformation retraction of  $Z/K$  to  $(\text{Crit}_G(V) \cap Z) / K$  for any  $G$ -invariant closed subset of  $V$ .
- The statement of the next result is simplified.
- Corollary.** If  $Z$  is Zariski closed in  $V$  then the GIT quotient,  $Z // G$ , of  $Z$  is a strict deformation retract of  $Z/K$ .
- This is a very useful result since if  $G$  is connected  $K$  is connected and this implies that  $Z // G$  has path lifting. In the complex case this is an important result of Kraft, Petrie and Randall.

- Let  $F(t, x)$  be the gradient flow corresponding to  $\phi$ . Then we have shown using freshman calculus that for  $t > 0$  and  $\|x\| \leq r$

$$\phi(F(t, x)) \leq \frac{C(r)}{t}.$$

- In addition if  $Z \subset V$  is closed and  $G$ -invariant then  $F(t, Z) \subset Z$  and 2 in the real Kempf-Ness theorem implies:
- Theorem.** Setting  $L(t, Kv) = KF(\frac{t}{1-t}, v)$   $0 \leq t < 1$  then  $\lim_{t \rightarrow 1} L(t, Kv)$  converges uniformly on compacta and this yields a strict deformation retraction of  $Z/K$  to  $(\text{Crit}_G(V) \cap Z) / K$  for any  $G$ -invariant closed subset of  $V$ .
- The statement of the next result is simplified.
- Corollary.** If  $Z$  is Zariski closed in  $V$  then the GIT quotient,  $Z // G$ , of  $Z$  is a strict deformation retract of  $Z/K$ .
- This is a very useful result since if  $G$  is connected  $K$  is connected and this implies that  $Z // G$  has path lifting. In the complex case this is an important result of Kraft, Petrie and Randall.
- We now consider the result implied by using the deep results of Lojasiewicz.

- The Lojasiewicz argument implies that if we set  $L(t, v) = F(\frac{t}{1-t}, v)$  then  $\lim_{t \rightarrow 1} H(t, v)$  converges uniformly on compacta.



- The Lojasiewicz argument implies that if we set  $L(t, v) = F(\frac{t}{1-t}, v)$  then  $\lim_{t \rightarrow 1} H(t, v)$  converges uniformly on compacta.
- **Theorem.** Let  $Z \subset V$  be closed and  $G$  invariant then  $L : [0, 1] \times Z \rightarrow Z$  defines a strong,  $K$ -equivariant deformation retraction of  $Z$  onto  $Z \cap \text{Crit}_G(V)$ .

- The Lojasiewicz argument implies that if we set  $L(t, v) = F(\frac{t}{1-t}, v)$  then  $\lim_{t \rightarrow 1} H(t, v)$  converges uniformly on compacta.
- **Theorem.** Let  $Z \subset V$  be closed and  $G$  invariant then  $L : [0, 1] \times Z \rightarrow Z$  defines a strong,  $K$ -equivariant deformation retraction of  $Z$  onto  $Z \cap \text{Crit}_G(V)$ .
- Over  $\mathbb{C}$  this result is due to Neeman.

- $\mathbb{C}^n = V \oplus iV$  so as a real vector space we write it as  $V \oplus V = \mathbb{R}^{2n}$ . The real part of the standard Hermitian inner product on  $\mathbb{C}^n$  becomes the standard inner product on  $\mathbb{R}^{2n}$ .  $M_n(\mathbb{C})$  becomes the algebra of  $2 \times 2$  block  $n \times n$  matrices

$$\begin{bmatrix} X & -Y \\ Y & X \end{bmatrix}.$$

Adjoint in  $M_n(\mathbb{C})$  becomes transpose in  $M_{2n}(\mathbb{R})$ .

- $\mathbb{C}^n = V \oplus iV$  so as a real vector space we write it as  $V \oplus V = \mathbb{R}^{2n}$ .  
The real part of the standard Hermitian inner product on  $\mathbb{C}^n$  becomes the standard inner product on  $\mathbb{R}^{2n}$ .  $M_n(\mathbb{C})$  becomes the algebra of  $2 \times 2$  block  $n \times n$  matrices

$$\begin{bmatrix} X & -Y \\ Y & X \end{bmatrix}.$$

Adjoint in  $M_n(\mathbb{C})$  becomes transpose in  $M_{2n}(\mathbb{R})$ .

- If  $X \subset \mathbb{C}^n$  is Zariski closed and defined by  $f_1, \dots, f_k$  in  $\mathbb{C}[x_1, \dots, x_n]$  then it is defined by  $\phi(x, y) = \sum |f_j(x + iy)|^2$  as a real variety.

- $\mathbb{C}^n = V \oplus iV$  so as a real vector space we write it as  $V \oplus V = \mathbb{R}^{2n}$ . The real part of the standard Hermitian inner product on  $\mathbb{C}^n$  becomes the standard inner product on  $\mathbb{R}^{2n}$ .  $M_n(\mathbb{C})$  becomes the algebra of  $2 \times 2$  block  $n \times n$  matrices

$$\begin{bmatrix} X & -Y \\ Y & X \end{bmatrix}.$$

Adjoint in  $M_n(\mathbb{C})$  becomes transpose in  $M_{2n}(\mathbb{R})$ .

- If  $X \subset \mathbb{C}^n$  is Zariski closed and defined by  $f_1, \dots, f_k$  in  $\mathbb{C}[x_1, \dots, x_n]$  then it is defined by  $\phi(x, y) = \sum |f_j(x + iy)|^2$  as a real variety.
- If  $G \subset GL(n, \mathbb{C})$  is a Zariski closed subgroup invariant under adjoint then  $G$  as a subgroup of  $GL(2n, \mathbb{R})$  is invariant under transpose. Furthermore, if we define the critical set for the action of  $G$  on  $\mathbb{C}^n$  to be

$$\{v \in \mathbb{C}^n \mid \langle Xv, v \rangle = 0, X \in \text{Lie}(G)\}$$

then this set is exactly  $\text{Crit}_G(\mathbb{R}^{2n})$ .

- $\mathbb{C}^n = V \oplus iV$  so as a real vector space we write it as  $V \oplus V = \mathbb{R}^{2n}$ . The real part of the standard Hermitian inner product on  $\mathbb{C}^n$  becomes the standard inner product on  $\mathbb{R}^{2n}$ .  $M_n(\mathbb{C})$  becomes the algebra of  $2 \times 2$  block  $n \times n$  matrices

$$\begin{bmatrix} X & -Y \\ Y & X \end{bmatrix}.$$

Adjoint in  $M_n(\mathbb{C})$  becomes transpose in  $M_{2n}(\mathbb{R})$ .

- If  $X \subset \mathbb{C}^n$  is Zariski closed and defined by  $f_1, \dots, f_k$  in  $\mathbb{C}[x_1, \dots, x_n]$  then it is defined by  $\phi(x, y) = \sum |f_j(x + iy)|^2$  as a real variety.
- If  $G \subset GL(n, \mathbb{C})$  is a Zariski closed subgroup invariant under adjoint then  $G$  as a subgroup of  $GL(2n, \mathbb{R})$  is invariant under transpose. Furthermore, if we define the critical set for the action of  $G$  on  $\mathbb{C}^n$  to be

$$\{v \in \mathbb{C}^n \mid \langle Xv, v \rangle = 0, X \in \text{Lie}(G)\}$$

then this set is exactly  $\text{Crit}_G(\mathbb{R}^{2n})$ .

- The original Kempf-Ness theorem is now a special case of the real Kempf-Ness theorem since Zariski closure of complex orbits is the same as the closure in the metric topology of  $\mathbb{R}^{2n}$ .

- The system in the abstract for my talk is just the case of  $GL(n, \mathbb{C})$  acting on  $M_n(\mathbb{C})$  by conjugation. Yielding the gradient system

$$\dot{X} = -4[[X, X^*], X].$$

- The system in the abstract for my talk is just the case of  $GL(n, \mathbb{C})$  acting on  $M_n(\mathbb{C})$  by conjugation. Yielding the gradient system

$$\dot{X} = -4[[X, X^*], X].$$

- Writing  $F_\infty(X) = \lim_{t \rightarrow +\infty} F(t, X)$  then  $F_\infty(X)$  is a normal operator with the same eigenvalues as  $X$ .







