

18.600 Midterm 1, Spring 2017: 50 minutes, 100 points

1. Carefully and clearly *show your work* on each problem (without writing anything that is technically not true). In particular, if you use any known facts (or facts proved in lecture) you should state clearly what fact you are using and why it applies.
2. No calculators, books, or notes may be used.
3. Simplify your answers as much as possible (but answers may include factorials — no need to multiply them out).

NAME: .....

1. (20 points) Let  $X$  be the number on a standard die roll (assuming values in  $\{1, 2, 3, 4, 5, 6\}$  with equal probability). Let  $Y$  and  $Z$  be the numbers on two independent rolls of the same die (so  $X$ ,  $Y$ , and  $Z$  are independent of each other). Compute the following:

(a)  $E[X]$  and  $E[X^2]$

(b)  $\text{Var}[X]$

(c)  $E[X^2 + 2Y^2 + 3Z^2]$

(d) The conditional probability  $P[X = 1|X = Y]$ .

2. (10 points) An urn contains 5 black balls and 5 white balls and 5 red balls and 5 yellow balls. If a collection of 8 balls is chosen uniformly at random from the urn, what is the probability that the collection contains exactly 2 balls of each color?

3. (20 points) Let  $M$  be the event that a person has a certain medical condition and let  $T$  be the event that a test of that condition comes back positive. Assume that the person has a  $1/20$  chance to have the condition and given that the person has the condition, there is a 90 percent chance the test will identify the condition correctly. Assume also that there is 20 percent chance the test will come back positive given that the person doesn't have the condition. In symbols, we can write the above assumptions as  $P(M) = 1/20$  and  $P(T|M) = 9/10$  and  $P(T|M^c) = 1/5$ . Now compute the following:

(a)  $P(T)$  (i.e., overall likelihood test is positive)

(b)  $P(M|T)$  (i.e., likelihood person has condition given test is positive)

(c)  $P(M|T^c)$  (i.e., likelihood person has condition given test is negative)

4. (10 points) A standard deck of cards has 4 suits, and 13 different cards of each suit (52 total). We know that there are  $\binom{52}{13}$  ways to choose an (unordered) set of 13 cards from this deck. How many ways are there to choose an (unordered) set of 13 cards with the property that 7 of the cards belong to the same suit (with the other 6 *not* belonging to that suit)?

5. (10 points) You have 12 pieces of pizza and 4 people at a party. How many ways are there to divide the 12 pieces among the four people? In other words, how many ordered four-tuples of non-negative integers  $(a_1, a_2, a_3, a_4)$  satisfy  $a_1 + a_2 + a_3 + a_4 = 12$ ?



7. (10 points) Fix an integer  $n \geq 3$ . Let  $\sigma$  be a random permutation chosen uniformly from the set of  $n!$  permutations of  $\{1, 2, \dots, n\}$ . Let  $X = X(\sigma)$  be the number of cycles of length 3 in the permutation  $\sigma$ . Compute  $E[X]$ .

Note: Here is some notation you can use in your answer if it is helpful.

Given integers  $i < j < k$ , let  $E_{i,j,k}$  be the event that the integers  $i, j, k$  are part of a length three cycle (which means that *EITHER*  $\sigma(i) = j$ ,  $\sigma(j) = k$ , and  $\sigma(k) = i$  *OR*  $\sigma(i) = k$ ,  $\sigma(k) = j$ , and  $\sigma(j) = i$ ). Let  $X_{i,j,k}$  be the indicator random variable  $1_{E_{i,j,k}}$ .